



TITLE:

Optimal culling strategy in relation to biological and economic efficiency and annualized net revenue in the Japanese Black cow-calf production system

AUTHOR(S):

OISHI, K.; IBI, T.; KAHI, A. K.; HIROOKA, H.

CITATION:

OISHI, K. ...[et al.]. Optimal culling strategy in relation to biological and economic efficiency and annualized net revenue in the Japanese Black cow-calf production system. *The Journal of Agricultural Science* 2011, 149(06): 783-799

ISSUE DATE:

2011-12

URL:

<http://hdl.handle.net/2433/150114>

RIGHT:

© Cambridge University Press 2011; この論文は出版社版ではありません。引用の際には出版社版をご確認ご利用ください。; This is not the published version. Please cite only the published version.

MODELLING ANIMAL SYSTEMS PAPER

Optimal culling strategy in relation to biological and economic efficiency and annualized net revenue in the Japanese Black cow-calf production system

Running head: Optimal culling strategy of beef cows in Japan

K. OISHI^{1*}, T. IBI², A. K. KAHI³ AND H. HIROOKA¹

¹ *Laboratory of Animal Husbandry Resources, Division of Applied Biosciences, Graduate School of Agriculture, Kyoto University, 606 8502 Kyoto, Japan*

² *Laboratory of Animal Breeding and Genetics, Division of Bioscience, Graduate School of Natural Science and Technology, Okayama University, 700 8530 Okayama, Japan*

³ *Animal Breeding and Genetics Group, Department of Animal Sciences, Egerton University, P. O. Box 536, 20115 Egerton, Kenya*

(Revised MS received 25 February 2011; Accepted 26 February 2011)

* To whom all correspondence should be addressed. Email: kazato@kais.kyoto-u.ac.jp

SUMMARY

The objective of the present study was to determine the optimal culling strategy in relation to biological and economic efficiency (BE and EE, respectively) and annualized net revenue (AN) in the Japanese Black cow-calf production system with special reference to the beef quality of culled cows. The herd model focused on two ways of mating: one-mating trial system (ONE) and continuous-mating trial system (CON). ONE assumed that heifers that fail to conceive are culled and cows that fail to conceive are culled at weaning of their calves while CON assumed that mating continues until all females theoretically conceive. Least squares means of carcass data of Japanese Black cows collected from a cooperative farm in Japan were used to estimate the carcass price of a cow by parity and Beef Marbling Standard (BMS) number. The simulation, assuming the current production situation in Japan, indicated that sales of culled cows accounted for 0.10–0.20 of total sales and was an important element in total production. Comparisons between ONE and CON showed that production efficiency in the current situation is higher in CON. The BE, EE and AN were higher in CON than in ONE. The two economic indicators were less sensitive to changes in annual discount rate but highly sensitive to changes in female calf price and BMS number of cows, indicating the importance of considering fluctuations in calf price and potential quality of culled cows' carcasses when estimating the economically optimal parity of culling. The three indicators derived different optimal solutions even in the same mating trial systems, stressing the importance of choice of production indicators when determining the culling strategy and evaluating animal production.

INTRODUCTION

Culling is important in managing cow-calf production systems because cow longevity and the carcass yield have major effects on profitability. Productivity in a cow-calf production system is influenced by cow age (Rogers 1972); however, beef quality and prices diminish with advancing age (Van Arendonk 1985). Therefore, profit may be lower if a cow is retained in the farm for too long.

The beef quality of a culled cow is evaluated in several countries using unique grading systems. In fact, returns from culled cows account for a significant part of the total returns of beef and dairy farms in Europe (Seegers *et al.* 1998; Garcia & Agabriel 2008) and sales of culled beef cows represent 0.10 to 0.20 of the gross revenue of cow-calf operations in the US (Sawyer *et al.* 2004). The strategies for culling cows may be affected by their beef quality traits (i.e. marbling). Hence, it is important to determine the effect of changes of beef quality in culled cows on the total production efficiency in different production systems.

Culling strategies in dairy cattle have been studied using numerous techniques including a representative dynamic programming model (Stewart *et al.* 1977; Van Arendonk 1985), dynamic programming model calculating the maximum average monthly return index (MaxAMR) (Kuipers 1982; Congleton & King 1985), hierarchic Markov process (Kristensen 1987), spreadsheet-based model (Meadows *et al.* 2005) and decision support systems to evaluate cull cow finishing strategies (Minchin *et al.* 2010). In beef cattle, Rogers (1972) investigated the optimal culling age of cows by including four variables: proportion of calf crop weaned, weaning weights of calves, sale value of cows and death loss of cows. Naazie *et al.* (1999) used a deterministic beef efficiency model to examine the effect of average culling age on overall life-cycle efficiency in the cow-calf production system. However, the effects of culling strategy on evaluation of total production by taking into account the change in beef quality and grading categories have been less widely studied.

Biological and economic evaluations have been conducted widely for several animal

production systems. Biological evaluations were carried out by calculating the biological efficiency (BE), determined as total production over total energy input (Cartwright 1979; Williams *et al.* 1995; Hirooka *et al.* 1998a) or vice versa; i.e. total energy input over total production (Tess *et al.* 1983). This measure, referred to as feed conversion ratio (Brody, 1945), is the most used index of production. The BE is stable because it is largely independent of changes in feed, calf and carcass prices resulting from economic fluctuations (Fowler *et al.* 1976). However, use of BE as a production indicator is limited because of lack of economic information (Dickerson 1970; Harris & Newman 1994; Hirooka *et al.* 1998b).

Economic evaluations have been done by calculating various economic values based on total revenue per unit of costs (Hirooka *et al.* 1998a), total cost per unit of product (Dickerson 1970), gross margin (Miller *et al.* 1999; Veyssset *et al.* 2005), the present value of profit per animal (Wolfová *et al.* 2009), etc. Economic efficiency (EE) is the total revenue per unit of costs or total cost per unit of product and is useful for evaluating economic situations in different planning horizon of production. Economic values such as gross margin and net present value are also widely used. They are simple and can be a production objective for producers, but they cannot be used for evaluation in different planning horizon of production. The annualized net revenue (AN) calculated from the net present value can solve the problem (Rogers *et al.* 1988). However, there are few studies that have investigated the effects of choice of method of biological and economic evaluations on culling strategies. The objective of the present study was to determine the optimal culling strategy in relation to BE, EE and AN in the Japanese Black cow-calf production system with special reference to the beef quality of culled cows.

MATERIALS AND METHODS

General

The model was developed to determine the optimal planned parity of culling of cows that

maximises the BE, EE and AN for a whole cycle in cow-calf production systems. The systems included birth, growth of calves in pre-weaning and post-weaning periods, selling male and non-replacement female calves at calf market, replacement of home-bred cows, repeated reproduction until the planned parity at culling, culling of cows after weaning of their calves and selling of culled cows' carcass. Production, reproduction and economic traits were incorporated into a herd composition model. Variables for nutrition and management were selected to represent specialized typical Japanese beef cow-calf production systems (NARO 2009). All simulations were conducted deterministically based on a one-day time step because production and reproduction traits in the standard are expressed on a daily basis. An explanation of the symbols and several fixed parameters used in the model are presented in Appendix 1.

Production traits

Daily mortality was considered for calves and cows. Body weight changes in each sex are estimated from growth curves. It was assumed that the growth curves were represented by a straight line from birth to weaning and by the Brody's curve (Brody 1945) from weaning to culling. The weaning age was assumed to be 150 days of age and is a fixed parameter. The weaning weight (WW) of calves was assumed to vary according to the changes in parity of their dams, and is expressed as a quadratic function as:

$$WW(n) = (-0.0091n^2 + 0.1081n + 0.7318) \times 180 \quad (1)$$

where n is the number of parities. The curving pattern of the change in weaning weight in Eqn (1), expressed by the quadratic function in parentheses, was derived from the data of Renquist *et al.* (2006). The function was multiplied by 180 in order to correct the expression to fit the situation in Japan (NARO 2009). Birth weight, weaning weight and mature weight of steers were expressed as those of heifers multiplied by 1.2, 1.08 and 1.2, respectively. For pregnant cows, total weight of the conceptus was added to maternal weight for the last 2 months of pregnancy (AFRC 1993). Daily milk yields were estimated using Wood's lactation curve

(Wood 1967). The daily metabolizable energy (ME) requirement was estimated based on AFRC (1993) and Japanese feeding standard for beef cattle (NARO 2009). The ME requirement of calves was calculated using the ratio of dam's milk to dietary feed on a dry matter (DM) basis. Details about estimation of mortality, body weight change, milk yields and ME requirement are shown in Appendix 2.

Reproduction traits

The conception rate of cows ($Cr(n)$) in each parity is calculated using a quadratic function estimated from Rogers *et al.* (1972) as:

$$Cr(n) = (-0.705n^2 + 6.264n + 80.159)/100 \quad (R^2 = 0.95) \quad (2)$$

where n indicates the number of parity. The $Cr(n)$ peaks in the 3rd parity and declines subsequently. The effect of differences in feeding level on the conception rate is not taken into account in the function, because the feed quantity is estimated from the ME requirement in the model and is assumed to be sufficient for mating. In the present study, two mating trial systems are assumed: a one-mating trial system (ONE) and Bailie's continuous-mating trials system (CON; Bailie 1982). The ONE assumes that heifers and cows are mated once in a breeding season and that heifers that fail to conceive are culled immediately and cows that fail to conceive are culled at weaning of their calves assuming a fixed calving interval of 357 days. In contrast, CON assumes that mating trial continues until all females theoretically conceive, in accordance with the procedure of Bailie (1982). In that procedure, the average period from parturition to the next conception for all females at parity n ($T_{do}(n)$, days) is determined by $Cr(n)$ assuming that the number of oestrous periods allowed for service is infinity. $T_{do}(n)$ is expressed as the sum of the anoestrus postpartum interval (t_{pp} , days) and the average of mating trial period for all females at parity n . The number of females that fail to conceive after i oestrous cycles decrease by $(1 - Cr(n))^i$. If mating continues until all females in parity n theoretically conceive, $T_{do}(n)$ is calculated as:

$$T_{do}(n) = (t_{pp} - t_{op}) + t_{op} \times Cr(n) + 2 \times t_{op} \times Cr(n)(1 - Cr(n)) \quad (3)$$

$$\begin{aligned}
 & + 3 \times t_{op} \times Cr(n)(1 - Cr(n))^2 + \dots \\
 & = (t_{pp} - t_{op}) + t_{op} \times \sum_{i=1}^{\infty} i(1 - Cr(n))^{i-1} \\
 & = (t_{pp} - t_{op}) + t_{op} / Cr(n)
 \end{aligned}$$

1 where t_{op} (days) is the mean length of the oestrous cycle and for first mating, and t_{pp} is
 2 equal to the sum of the weaning and post-weaning periods; i.e. the age at first mating (t_{mfst} ,
 3 days). It is then assumed that all females in parity n conceive on the same day represented
 4 by $T_{do}(n)$ and that only cows with the planned parity at culling are culled at weaning of their
 5 calves. The average number of mating trial times ($Mt(n)$, n) for all females in parity n in
 6 the herd is then calculated as:

$$\begin{aligned}
 Mt(n) &= 1 + (1 - Cr(n)) + (1 - Cr(n))^2 + (1 - Cr(n))^3 + \dots \\
 &= \sum_{i=1}^{\infty} (1 - Cr(n))^{i-1} \\
 &= 1 / Cr(n).
 \end{aligned} \tag{4}$$

7 Finally, the calving interval $T_{cl}(n)$ (days) is defined as the period between parturitions and is
 8 the sum of $T_{do}(n)$ and the gestation length (t_{preg} , days) as:

$$T_{cl}(n) = T_{do}(n) + t_{preg}. \tag{5}$$

9 In the present study, the conception rate was treated as the relative number of calves produced
 10 without considering prenatal calf mortality, to simplify the comparison between the two
 11 mating trial systems, and t_{op} , t_{mfst} and t_{preg} were treated as fixed parameters and were set
 12 to 21, 420 and 285 days, respectively.

13

14 *Determination of cow carcass price and calf price*

15 Carcass data of culled cows were obtained from farms consigned under the Agura Farm
 16 Enterprise, which is the largest cooperative farming company for Japanese Black cattle in Japan
 17 (Ibi *et al.* 2005). The data comprised 9759 records of culling date, carcass market, slaughter age
 18 (days), parity (n), fattening period (days), body weight (kg), carcass weight (kg), marbling

score (n), and carcass price (¥/kg) of culled cows. The marbling score is measured according to the Beef Marbling Standard (BMS), with scores, also known as BMS number, of 1 to 12 where 12 is the best (JMGA 1988). Note that in Japan, carcass price is determined at auction and reflects the quality of carcasses from culled cows as well as feedlot animals.

Table 1 shows the least squares means of carcass prices (CP , ¥/kg carcass) by culling parity and BMS number, estimated by adjusting for the effects of slaughter year and month and carcass market using the PROC GLM of SAS (1999). The effect of fattening period was excluded from the adjusting procedure because its effect on carcass price was not significant. Culling age was also excluded from the procedure since age was strongly correlated with culling parity in the data. The quadratic regressions shown in Fig. 1 were obtained using PROC REG of SAS (1999) and the equations to estimate CP (¥/kg carcass) based on culling parity and BMS number are as follows:

$$\begin{aligned} CP(pa) &= -1.0038pa^2 + 2.875pa + 589.3 \quad (R^2 = 0.52) \quad (BMS = 1) \\ CP(pa) &= 0.2559pa^2 - 22.571pa + 909.8 \quad (R^2 = 0.93) \quad (BMS = 2) \\ CP(pa) &= 1.7145pa^2 - 42.475pa + 1118.6 \quad (R^2 = 0.94) \quad (BMS = 3) \\ CP(pa) &= 3.0998pa^2 - 70.304pa + 1397.1 \quad (R^2 = 0.88) \quad (BMS \geq 4). \end{aligned} \quad (6)$$

where pa is parity of cows at culling. Dressing percentage for calculating carcass weight of culled cows was set to 61.35%, which is the average value of carcass weight per body weight in the carcass data.

The default female calf price of ¥1250 per live weight kg was assumed based on average data for the last 10 years (MAFF 2008). Relative calf price ratio of live male to live female was fixed and set to 1.1344.

Setting of cost parameters

For simplicity, only two categories of feeds are assumed in the present study: purchased roughage and concentrates. Seasonal changes in feed ingredients were not taken into account. Feed costs were calculated by multiplying feed prices for different animal classes using the feed

requirement. For pre-weaning calves, fixed parameters of efficiency of ME utilization of milk and dietary feed were assumed to estimate total feed requirement. The ratio of concentrates to the dietary feed for pre-weaning calves was set as an age-related linear function derived from JLTA (2007). For post-weaning steers, heifers and cows, feed requirements from roughage and concentrates were estimated based on metabolizability as described by Hirooka *et al.* (1998). Details about calculation of feed amounts for each animal category are shown in Appendix 2. Prices of concentrates were assumed to be ¥30 and ¥38/ kg DM for cows and pre-weaning calves, respectively. Price of roughage was assumed to be 1.30 times higher than of concentrates according to the situation in Japan. As for other production costs, most beef cows are artificially inseminated (AI) in Japan and the cost of AI per oestrous cycle was assumed to be ¥12000. The average cost of AI for all females at parity n were then calculated as the cost of AI per oestrous period multiplied by the average number of mating trials for all females at parity n ($Mt(n)$). The other costs, including managerial costs and machinery costs, was assumed to total ¥392 per day for producing a calf (MAFF 2009).

Herd composition dynamics

Figure 2 shows the outline of the herd composition dynamics. Three animal categories can be distinguished: male calves, non-replacement female calves, and cows. Although the number of cows decreases between each reproduction because of the change in survival rate and the culling of non-pregnant cows in ONE, it is assumed that subsequent home-bred replacement heifers join the herd in each reproduction. It was assumed that all male calves and non-replacement female calves after weaning are sold at 285 days of age in calf markets after post-weaning growing period. In the present study, optimal culling parity was simulated between parities 3-12. This was because the calculated result using the herd composition dynamics model indicated that the herd cannot maintain the initial number of cows with home-bred replacement heifers only when reproduction is less than three times. There were no carcass data for culled cows with $BMS \geq 4$ over 12th parity. Cows were culled when their

calves born in the optimal culling parity were weaned. Fattening period of cows was not considered in the model because it did not significantly affect carcass price of cows in the GLM analysis described above. Total herd outputs were calculated as individual production traits multiplied by animal numbers in herd components derived from herd composition dynamics. Details of the herd composition dynamics are shown in Appendix 3.

Net present value and annualized net revenue

The production objective in most enterprises is to maximize the net present value (NPV) of the entire future stream of residual earnings from the productive process (Perrin 1972), and cull and replace cows when it is optimal to maximize this present value. The NPV of a series of future cash flows, so-called the discounted cash flow, is calculated as follows:

$$NPV(pa) = \sum_{i=1}^{Day(pa)} (CF(i) / (1 + dr)^i) \quad (7)$$

where $NPV(pa)$ is the net present value associated with keeping cows from birth to culling when planned culling parity is set to pa , $Day(pa)$ is the planning days of age of cows when planned culling parity is set to pa , $CF(i)$ is the daily cash flow and dr is the daily discount rate. The daily cash flow is defined as the daily return (estimated only in cases where calves or beef from culled cows are sold) minus daily cost (including feed cost, AI cost and other fixed cost) for cows with i age in days and their calves. The daily discount rate is calculated from annual discount rate as: $dr = \sqrt[365]{(1 + ydr)} - 1$ where ydr is the annual discount rate. Here, the discount rate includes three components: a risk-free rate for time preference, an inflation premium, and a risk premium (Barry *et al.* 1995).

However, NPV comparisons are strictly limited to comparison of replaceable assets having equal life spans (Meadows *et al.* 2005). Cows kept for long periods should have a higher NPV as there is more production relative to the cows' feeding cost. To eliminate this time problem, the NPV associated with each planned culling needs to be converted to an equivalent yearly annuity. This equivalent yearly annuity would be the constant amount of net revenue each

1 year available to farmers (Rogers *et al.* 1988), and is expressed using the following formula:

$$\begin{aligned} EDC(pa) &= NPV(pa) \times dr / (1 - (1/(1 + dr))^{Day(pa)}) \\ AN(pa) &= EDC(pa) \times 365 \end{aligned} \quad (8)$$

2 where $EDC(pa)$ is the equivalent daily cash flow associated with keeping cows from birth
3 to culling when planned culling parity is set to pa , and $AN(pa)$ is the annualized net
4 revenue (AN) (or the estimated equivalent annuity) using $EDC(pa)$ (Meadows *et al.* 2005).
5 The planned parity at culling with the highest AN represents the optimal targeted herd life.

7 *Evaluation of model outputs*

8 The model developed in the present study evaluates AN and two other production indicators:
9 BE and EE. The BE was defined as total sale live weights of a culled cow and her calves per
10 unit of total ME intake of the herd from birth to culling. The EE was determined as returns over
11 costs, including feed costs and other production costs from birth to culling.

12 The simulation was first performed under a base situation of Japanese cow-calf production in
13 two mating trial systems. In the base situation, culling parity of cows, female calf price,
14 Japanese Beef Marbling Standard number and annual discount rate were assumed to be 6,
15 ¥1250, 3 and 5%, respectively. Sensitivity of BE, EE and AN to changes in conception rate,
16 weaning weight, weaning age, calf market age in both mating trial system was performed. In
17 addition, sensitivity of EE and AN to changes in female calf price and annual discount rate was
18 also performed. For the sensitivity analysis, fixed weaning weight ($WW = 180$) and fixed
19 conception rate ($Cr = 0.8$) were used instead of the weaning weight change function described
20 in Eqn (1) and the conception rate change function described in Eqn (2). The relationships
21 between optimal culling parity and the three indicators were investigated. In addition,
22 difference between the effects of the two economic indicators on determination of optimal
23 culling strategies was analysed.

25 **RESULTS**

Simulated outputs under the base situation

Table 2 shows the simulated outputs per cow under the base situation of Japanese cow-calf production in the two mating trial systems. The Japanese cow-calf production system has so far focused only on revenue from sale calves. However, the sales of culled cows accounted for 17.5% in ONE and 11.8% in CON. Comparisons between the two mating trial systems showed that the number of newborn calves, BE, EE and AN were all higher in CON than in ONE.

Sensitivity of biological and economic efficiency and annualized net revenue

Figure 3 shows the sensitivity of BE, EE and AN to changes in conception rate, weaning weight, weaning age, calf market age, female calf price and annual discount rate in the two mating trial systems. The three indicators were all sensitive to changes in parameters, but the level of sensitivity was different among them. The AN was the most and the BE the least sensitive to changes in the parameters. The three indicators were highly sensitive to changes in conception rate in ONE than in CON. The sensitivity of the indicators to changes in the other parameters was higher in CON than in ONE but the differences in the level of sensitivities between the two mating trial systems were small. The influence of female calf price on the two economic indicators was large but that of annual discount rate was small.

Effect of culling parity on biological efficiency

The effect of changes in culling parity of cows on BE in the two mating trial systems is presented in Fig. 4. An increase in culling parity led to a decrease in BE in both systems. When comparing the two mating trial systems, BE in CON was higher than in ONE in all parities of culling but the pattern of decrease was different between the two systems. In ONE, the pattern of the decrease mainly resulted from the changes in conception rate and weaning weight of calves. In addition to the two parameters, the pattern of decrease in CON was greatly affected by the interaction between the reduction and improvement in BE which were caused by

extension of mating trial period and constant production of calves having high efficiency of feed utilisation.

Effect of culling parity, BMS number and female calf price on economic efficiency and annualized net revenue

Figure 5 shows the effects of culling parity, BMS number, female calf price and annual discount rate on EE in the two mating trial systems. The optimal culling parity varied depending on levels of BMS number and the female calf price but did not vary greatly with levels of annual discount rate. Annual discount rate only negatively affected the magnitude of the EE in both systems. When female calf price was ¥1000, optimal culling parity decreased with an increase in BMS number in both systems. However, optimal culling parity gradually increased until the 9th parity in all BMS number categories when female calf price was raised to ¥1500. Decrease in EE in later parity of culling was a result of the decrease in conception rate and weaning weight of calves. Values of EE were higher in CON than in ONE, especially when female calf price was high and annual discount rate low.

Figure 6 shows the effects of culling parity, BMS number, female calf price and annual discount rate on AN in the two mating trial systems. As was the case for EE, the optimal culling parity varied depending on levels of BMS number and the female calf price. The effect of the change in annual discount rate on AN was slightly larger than on EE. Levels of annual discount rate negatively affected values of AN in both systems, in a similar fashion to EE. When female calf price was ¥1500, AN decreased substantially in later parity of culling in ONE whereas the rate of the decrease was very low in CON. The decrease in AN in ONE in later parities was associated with a decline in the conception rate because this strongly affected AN in ONE as shown in Fig. 3. Values of AN were higher in CON than in ONE; consistent with the findings for EE.

The differences in optimal culling parity between EE and AN with changes in BMS number, female calf price, annual discount rate and the mating trial system, are presented in Fig. 7. This

clarifies the difference between EE and AN. When female calf price was set to ¥1000, both indicators in both mating trial systems derived same optimal solutions for BMS number 4 indicating that culling early is the best. For other price settings, however, the two indicators in the two systems did not show same optimal culling parities. An important finding was that EE and AN derived different optimal solutions even in the same mating trial systems. This difference was particularly prominent in the highest BMS numbers with female calf price of ¥1250 in CON. The optimal solutions of AN appear to be relatively stable especially when female calf price was equal to or higher than ¥1250.

Break-even values of female calf price

Table 3 shows the break-even values of female calf price when the AN in ONE exceeds the revenue in CON with culling in 6th parity. These values indicate that ONE should be selected when calf price is lower than the break-even values. The break-even values clearly increase consistently with annual discount rate, and decrease with an increase in conception rate. The effect of the change in BMS number on the break-even value is negative when conception rate is ≤ 0.40 and positive when conception rate is ≥ 0.80 . The effect of the change in conception rate on the break-even value was the largest among the three parameters.

Table 4 shows the break-even values of female calf price when AN at culling in the 9th parity exceeds the revenue at culling in the 3rd parity for CON. These were calculated to verify the effect of changes in calf price, BMS number and annual discount rate on selection of early culling (3rd parity) or late (9th parity). Here, late culling was assumed to be in 9th parity, because the average value of optimal culling parity estimated by AN in CON (except when the optimal solutions indicated that the earliest culling was optimal) was equal to 9. The break-even values clearly increase with an increase in BMS number and annual discount rate. The change in annual discount rate was linear to the change in the break-even value. The effect of the change in BMS number on the break-even value was larger than in annual discount rate.

DISCUSSION

Evaluation of the base situation

Under the base situation in Japanese cow-calf production, the sales of culled cows accounted for over 10% in both mating trial systems. This is consistent to the report of Sawyer *et al.* (2004) in US representing 10 to 20% of contribution of culled cows' beef to the gross revenue. The present result indicated that culled cows are also important revenue sources in Japan. In addition, BE, EE and AN were all higher in CON than in ONE under the base situation. Although the mating cost increases with an increase in mating trial times and the sales of cow carcasses decrease with advancing age at culling, the result indicated that the benefit of calf production exceeded the cost and the loss in sales in the base situation.

Sensitivity of the three production indicators

In the sensitivity analyses, the AN was the most and the BE the least sensitive to changes in the parameters and the three indicators were highly sensitive to changes in conception rate in ONE than in CON. The difference in sensitivity indicates the high dependency of conception rate in ONE since failure to mate will directly affect the production outputs.

The negative influence of annual discount rate on the two economic indicators was smaller than expected. This may be due to the small changes in the rate in the present study. For example, a 20% change in the annual discount rate of 5% is equivalent to only a 1% change in the discount rate. On the other hand, the two economic indicators were highly sensitive to changes in female calf price. This result indicated the importance of considering fluctuations in calf price when estimating the economically optimal culling age of cows, consistent with the study of Clarke *et al.* (1984a).

Relationships between culling parity and the three indicators

BE decreased consistently with increasing culling parity in both systems. Similar results have

been reported elsewhere (Taylor *et al.* 1985; Bourdon & Brinks 1987; Baptist 1992). Taylor *et al.* (1985) reported a decline in maximum overall efficiency of food utilisation as the number of calving per dam increased. Baptist (1992) also reported that an increase in culling age of cows resulted in a decrease in BE.

In contrast, optimal culling parity greatly varied with the changes in economic parameters when EE and AN were used as production indicators. When female calf price was ¥1000, the two economic indicators were maximized at the lowest culling parity for the highest BMS number and at later parity of culling for the lowest BMS number. Reduction in female calf price led to a low optimal culling parity, especially when BMS number was high. At low calf prices, revenue from sales of culled cows contributed greatly to the total sales because of low sales of its calves. This clarified the effect of differences in carcass quality of cows on the economic indicators.

Comparison of three production indicators

The patterns of change in BE and AN were similar when female calf price was lower than the average price of culled cows for each BMS number. The decrease in BE in the present study appears to be due to the fact that the difference in prices between culled cows and their calves was not considered in calculating BE. The change in BE corresponds to the extreme case of no price difference between culled cows and sales of calves in AN. Hence, AN can express production efficiency just like BE according to the changes in setting of production circumstances. Comparison between the use of EE and AN in determining optimal parity at culling in this study indicates that choice of economic indicators is important for evaluating animal production systems. Sensitivity of biological and economic parameters on AN was higher than on EE. This suggests that AN is more effective in the evaluation of animal productions than EE. The AN directly implies production benefit and thereby it is easily understandable by decision makers. Therefore, use of the AN is recommended as a measure of production efficiency.

Selections of ONE or CON and early or late culling

The results in Table 3 indicated that CON is better than ONE unless calf price is extremely low. Even though extension of calving period by a decrease in conception rate accumulates costs of mating and feeding and also decreases the quality of cow carcasses, increasing sales of calves by continuous mating can compensate for the negative effects of CON. Therefore mating should continue insofar as breeding season continues. Clarke *et al.* (1984b) simulated culling strategies for beef production and concluded that culling non-pregnant cows improved EE, however the relative price of live calves to non-pregnant cows was low in that study. Hence, relatively higher calf price compared with culled cow carcass price should give high NPV for calf production.

The mean prices of the break-even values of female calf price in Table 4 were all higher than the average carcass prices of culled cows with corresponding BMS number calculated from Table 1 (BMS number 1: ¥556, 2: ¥787, 3: ¥950, 4: ¥1130). This result indicated that culling in early parity is the best choice when female calf price is the same as carcass price of culled cows.

The mean prices in Table 3 and Table 4 indicated that, in the actual Japanese Black cow-calf production, cows should be mated continuously to the extent possible and be culled in later parities, because the average price of live calves for the last 10 years in Japan has been ¥1250, which was higher than the mean values.

Effect of carcass price of culled cows

In general, culling of cows later in their lifetime may be caused by high sales price of calves compared to that of culled dams. For commercial beef cow-calf production in USA, Melton *et al.* (1994) developed a bio-economic simulation model and reported that an increase in culled cow price relative to the calf price results in cows being culled at an earlier age. Naazie *et al.* (1997) mentioned that the decline in overall efficiency with an increase in culling age of cows

may be a result of high price of culled cows. Bourdon & Brinks (1987) concluded that optimal age at culling of cows has a negative relationship with relative price of culled cow to her calves. The present studies are consistent with those results, indicating that calf prices lower than the break-even values (Table 4) resulted in the reduction in optimal culling parity.

The calf market situation in Japan is dynamic. The average calf price per kg body weight went below ¥1000 when cases of bovine spongiform encephalopathy (BSE) were reported in Japan (MAFF 2008). The present study showed that, when the price of live calves is low, the difference in BMS number of cows should be of great importance in determining the economically optimal culling parity of cows. This result stresses the importance of the difference in beef quality of culled cows in optimization of whole production, and thus it may be necessary to predict BMS number of cows before effecting culling decisions.

The differences in beef quality have raised concern when evaluating cattle production systems not only in Japan but also in Europe and the US. Bekman & Van Arendonk (1993) introduced the EUROP classification into their model to estimate economic values for veal, beef and milk production traits and evaluate the difference in the quality of beef and veal. Linamo & Van Arendonk (1999) also analysed the effect of changes in price difference between carcass categories of dairy cows on predicted total monetary response in alternative selection strategies. In the US, a relationship between carcass value and body condition of beef cows to optimize economic returns and USDA carcass grading information has been reported (Apple 1999). Beef quality is evaluated primarily by physical characteristics that are associated with age (Yager *et al.* 1980) or reproduction, as shown in the present study. Therefore, culling policy should be examined in the context of differences in beef grade of culled cows.

CONCLUSIONS

The present study showed the differences between biological efficiency, economic efficiency and annualized net revenue in determining the optimal culling strategy. Comparisons between

the three indicators showed that they all derived different optimal solutions even in the same mating trial systems, stressing the importance of choice of production indicators when determining the culling strategy and evaluating animal production. In addition, the study demonstrated the importance of culled cows' beef quality when determining economically optimal culling parity of cows in the Japanese Black cow-calf production system. The changes in optimal culling parity and the price changes of culled cows' beef as a result of the differences in beef grade play a significant role in the economic evaluation of beef production systems.

REFERENCES

- AGRICULTURAL AND FOOD RESEARCH COUNCIL (AFRC). (1993). *Energy and Protein Requirements of Ruminants*. An advisory manual prepared by the Agricultural and Food Research Council Technical Committee on Responses to Nutrients. CAB International, Wallingford.
- APPLE, J.K. (1999). Influence of body condition score on live and carcass value of cull beef cows. *Journal of Animal Science* **77**, 2610-2620.
- BAILIE, J.H. (1982). The influence of breeding management efficiency on dairy herd performance. *Animal Production* **34**, 315-323.
- BAPTIST, R. (1992). Derivation of steady-state herd productivity. *Agricultural Systems* **39**, 253-272.
- BARRY, P.J., ELLINGER, P.N., HOPKIN, J.A. & BAKER, C.B. (1995). *Financial Management in Agriculture*, 5th edn. Interstate Publishers Inc., Danville, Illinois.
- BEKMAN, H. & VAN ARENDONK, J.A.M. (1993). Derivation of economic values for veal, beef and milk production traits using profit equations. *Livestock Production Science* **34**, 35-56.
- BOLEMAN, S.J., BOLEMAN, S.L., MILLER, R.K., TAYLOR, J.F., CROSS, H.R., WHEELER, T.L., KOOHARAIE, M., SHACKELFORD, S.D., MILLER, M.F., WEST, R.L., JOHNSON, D.D. & SAVELL, J.W. (1997). Consumer evaluation of beef of known categories of tenderness.

- 1 *Journal of Animal Science* **75**, 1521-1524.
- 2 BOURDON, R.M. & BRINKS, J.S. (1987). Simulated efficiency of range beef production. III.
- 3 Culling strategies and nontraditional management systems. *Journal of Animal Science* **65**,
- 4 963-969.
- 5 BRODY, S. 1945. *Bioenergetics and Growth*. Reinhold Publishing Corporation, New York.
- 6 CARTWRIGHT, T.C. (1979). Size as a component of beef production efficiency: Cow-calf
- 7 production. *Journal of Animal Science* **48**, 974-980.
- 8 CLARKE, S.E., GASKINS, C.T., HILLERS, J.K. & HOHENBOKEN, W.D. (1984a). Mathematical
- 9 modeling of alternative culling and crossbreeding strategies in beef production. *Journal of*
- 10 *Animal Science* **58**, 6-14.
- 11 CLARKE, S.E., GASKINS, C.T., HILLERS, J.K. & HOHENBOKEN, W.D. (1984b). Mathematical
- 12 modeling of alternative selection strategies for beef production. *Journal of Animal Science* **59**,
- 13 308-316.
- 14 CONGLETON, W.R., Jr. & KING, L.W. (1985). Culling cows on predicted future profitability with
- 15 a variable planning horizon. *Journal of Dairy Science* **68**, 2970-2983.
- 16 DICKERSON, G.E. (1970). Efficiency of animal production – Molding the biological component.
- 17 *Journal of Animal Science* **30**, 849-859.
- 18 FERRIER, P. & LAMB, R. (2007). Government regulation and quality in the US beef market.
- 19 *Food Policy* **32**, 84-97.
- 20 FOWLER, V.R., BICHARD, M. & PEASE, A. (1976). Objectives in pig breeding. *Animal*
- 21 *Production* **23**, 365-387.
- 22 GARCIA, F. & AGABRIEL, J. (2008). CompoCow: a predictive model to estimate variations in
- 23 body condition and the energy requirements of culled cows during finishing. *Journal of*
- 24 *Agricultural Science, Cambridge* **146**, 251-265.
- 25 HARRIS, D.L. & NEWMAN, S. (1994). Breeding for profit: synergism between genetic
- 26 improvement and livestock production (a review). *Journal of Animal Science* **72**, 2178-2200.
- 27 HIROOKA, H., GROEN, A.F. & HILLERS, J. (1998a). Developing breeding objectives for beef

- 1 cattle production. 1. A bio-economic simulation model. *Animal Science* **66**, 607-621.
- 2 HIROOKA, H., GROEN, A.F. & HILLERS, J. (1998*b*). Developing breeding objectives for beef
- 3 cattle production. 2. Biological and economic values of growth and carcass traits in Japan.
- 4 *Animal Science* **66**, 623-633.
- 5 IBI, T., HIROOKA, H., KAHI, A.K., SASAE, Y. & SASAKI, Y. (2005). Genotype × environment
- 6 interaction effects on carcass traits in Japanese Black cattle. *Journal of Animal Science* **83**,
- 7 1503-1510.
- 8 JAPANESE MEAT GRADING ASSOCIATION (JMGA). (1998). *New Beef Grading Standards*.
- 9 Japanese Meat Grading Association, Tokyo. (In Japanese)
- 10 JAPAN LIVESTOCK TECHNOLOGY ASSOCIATION (JLTA). (2007). *Manual To Prevent Loss of*
- 11 *Japanese Black Calves*. Japan Livestock Technology Association, Tokyo. (In Japanese)
- 12 KRISTENSEN, A.R. (1987). Optimal replacement and ranking of dairy cows determined by a
- 13 hierarchic Markov process. *Livestock Production Science* **16**, 131-144.
- 14 KUIPERS, A. (1982). Development and economic comparison of selection criteria for cows and
- 15 bulls with a dairy herd simulation model. Ph.D. thesis. Agricultural Research Reports 913,
- 16 Centre for Agricultural Publishing and Documentation, Wageningen.
- 17 LINAMO, A.E. & VAN ARENDONK, J.A.M. (1999). Combining selection for carcass quality,
- 18 body weight, and milk traits in dairy cattle. *Journal of Dairy Science* **82**, 802-809.
- 19 MEADOWS, C., RAJALA-SCHULTZ, P.J. & FRAZER, G.S. (2005). A spreadsheet-based model
- 20 demonstrating the nonuniform economic effects of varying reproductive performance in
- 21 Ohio dairy herds. *Journal of Dairy Science* **88**, 1244-1254.
- 22 MELTON, B.E., ARDEN COLETTE, W., SMITH, K. & WILLHAM, R.L. (1994). A time-dependent
- 23 bioeconomic model of commercial beef breed choices. *Agricultural Systems* **45**, 331-347.
- 24 MILLER, S.P., WILTON, J.W. & PFEIFFER, W.C. (1999). Effects of milk yield on biological
- 25 efficiency and profit of beef production from birth to slaughter. *Journal of Animal Science* **77**,
- 26 344-352.
- 27 MINCHIN, W., O'DONOVAN, M., BUCKLEY, F. KENNY, D.A. & SHALLOO, L. (2010).

- 1 Development of a decision support tool to evaluate the financial implications of cull cow
- 2 finishing under different feeding strategies. *Journal of Agricultural Science, Cambridge*
- 3 Published online by Cambridge University Press 08 Apr 2010
- 4 MINISTRY OF AGRICULTURE, FORESTRY AND FISHERIES OF JAPAN (MAFF). (2008). *The change*
- 5 *in average price of beef calves*. Livestock Industry Department, Agricultural Production
- 6 Bureau, Ministry of Agriculture, Forestry and Fisheries of Japan, Tokyo. (In Japanese)
- 7 MINISTRY OF AGRICULTURE, FORESTRY AND FISHERIES OF JAPAN (MAFF). (2009). *Statistics of*
- 8 *agriculture, forestry and fisheries*. Statistics division of the Ministry of Agriculture, Forestry
- 9 and Fisheries of Japan, Tokyo. (In Japanese)
- 10 NAAZIE, A., MAKARECHIAN, M. & HUDSON, R.J. (1997). Efficiency of beef production
- 11 systems: description and preliminary evaluation of a model. *Agricultural Systems* **54**,
- 12 357-380.
- 13 NAAZIE, A., MAKARECHIAN, M. & HUDSON, R.J. (1999). Evaluation of life-cycle herd
- 14 efficiency in cow-calf systems of beef production. *Journal of Animal Science* **77**, 1-11.
- 15 NATIONAL AGRICULTURE AND FOOD RESEARCH ORGANIZATION (NARO). (2009). *Japanese*
- 16 *feeding standard for beef cattle*. National Agriculture and Food Research Organization,
- 17 Central Association of Livestock Industry, Tokyo. (In Japanese)
- 18 PERRIN, R.K. (1972). Asset replacement principles. *American Journal of Agricultural*
- 19 *Economics* **54**, 60-67.
- 20 RENQUIST, B.J., OLTJEN, J.W., SAINZ, R.D. & CALVERT, C.C. (2006). Effects of age on body
- 21 condition and production parameters of multiparous beef cows. *Journal of Animal Science* **84**,
- 22 1890-1895.
- 23 ROGERS, LEROY F. (1972). Economics of replacement rates in commercial beef herds. *Journal*
- 24 *of Animal Science* **34**, 921-925.
- 25 ROGERS, G.W., VAN ARENDONK, J.A.M. & MCDANIEL, B.T. (1988). Influence of production
- 26 and prices on optimum culling rates and annualized net revenue. *Journal of Dairy Science* **71**,
- 27 3453-3462.

- 1 SAS. (1999). *SAS/STAT User's Guide. Version 8*. SAS Institute Inc., Cary, NC, USA.
- 2 SAWYER, J.E., MATHIS, C.P. & DAVIS, B. (2004). Effects of feeding strategy and age on live
3 animal performance, carcass characteristics, and economics of short-term feeding programs
4 for culled beef cows. *Journal of Animal Science* **82**, 3646-3653.
- 5 SEEGER, H., BAREILLE, N. & BEAUDEAU, F. (1998). Effects of parity, stage of lactation and
6 culling reason on the commercial carcass weight of French Holstein cows. *Livestock*
7 *Production Science* **56**, 79-88.
- 8 STEWART, H.M., BURNSIDE, B.E. & WILTON, J.W. (1977). A dynamic programming approach to
9 culling decisions in commercial dairy herds. *Journal of Dairy Science* **60**, 602-617.
- 10 TAYLOR, S.T., C.S., MOORE, J.A., THIESSEN, R.B. & BAILEY, C.M. (1985). Efficiency of food
11 utilization in traditional and sex-controlled systems of beef production. *Animal Production*
12 **40**, 401-440.
- 13 TESS, M.W., BENNETT, G.L. & DICKERSON, G.E. (1983). Simulation of Genetic Changes in Life
14 Cycle Efficiency of Pork Production. I. A Bioeconomic Model. *Journal of Animal Science* **56**,
15 336-353.
- 16 VAN ARENDONK, J.A.M. (1985). Studies on the replacement policies in dairy cattle. II.
17 Optimum policy and influence of changes in production and prices. *Livestock Production*
18 *Science* **13**, 101-121.
- 19 VEYSSET, P., BEBIN, D. & LHERM, M. (2005). Adaptation to Agenda 2000 (CAP reform) and
20 optimisation of the farming system of French suckler cattle farms in the Charolais area: a
21 model-based study. *Agricultural Systems* **83**, 179-202.
- 22 WILLIAMS, C.B., BENNETT, G.L. & KEELE, J.W. (1995). Simulated influence of postweaning
23 production system on performance of different biological types of cattle: III. Biological
24 efficiency. *Journal of Animal Science* **73**, 686-698.
- 25 WOLFOVÁ, M., WOLF, J., KRUPOVÁ, Z. & KICA, J. (2009). Estimation of economic values for
26 traits of dairy sheep: I. Model development. *Journal of Dairy Science* **92**, 2183-2194.
- 27 WOOD, P.D.P. (1967). Algebraic model of the lactation in cattle. *Nature* **216**, 164-165.

- 1 YAGER, W.A., GREER, R.C. & BURT, O.R. (1980). Optimal policies for marketing cull beef
- 2 cows. *American Journal of Agricultural Economics* **62**, 456-467.

1

Table 1. *Least squares means of carcass price (CP*) by BMS[†] for Japanese Black cows*

Parity	BMS=1		BMS=2		BMS=3		BMS≥4	
	No.	CP	No.	CP	No.	CP	No.	CP
0	8	615	431	944	456	1169	363	1506
1	9	600	485	879	469	1054	209	1271
2	6	532	319	844	329	1010	141	1205
3	17	610	434	842	347	992	205	1166
4	19	617	444	799	354	969	184	1143
5	27	515	525	793	373	940	155	1138
6	51	549	573	778	337	929	92	1098
7	50	567	434	780	240	912	51	1126
8	34	599	377	768	146	913	41	1049
9	33	565	200	743	89	884	15	1006
10	15	494	146	719	46	884	10	1013
11	17	523	129	709	36	860	8	1010
12	31	447	170	641	66	826	13	961

* Least squares means of carcass price (yen/kg carcass weight).

[†] Japanese Beef Marbling Standard number.

2

1

Table 2. *Simulated outputs per cow under the base situation**

Outputs per cow	Mating system [†]	
	ONE	CON
No. of calves born	3.832	5.424
Biological output		
Live weight (kg)		
Culled cow	406	439
Live calves	806	1258
Total	1211	1696
Food intake (GJ)		
Culled cow	124.88	154.36
Live calves	41.27	63.62
Total	166.15	218.01
Total biological efficiency (kg/GJ)	7.29	7.78
Economic output (including the discount effect)		
Sales (yen)		
Culled cow	184033	176938
Live calves	864654	1322682
Total	1048688	1499620
Cost (yen)		
Food cost		
Culled cow	357366	447608
Live calves	120044	182805
Total	477410	630413
AI cost	59846	65117
Other costs	192961	297625
Total	730217	993155
Total benefit (yen)	318471	506465
Total economic efficiency	1.436	1.510
Annualized net revenue (yen)	54502	86451

* Culling parity of cows, female calf price, Beef Marbling Standard number and annual discount rate were set to 6, 1250, 3 and 5%, respectively.

[†]One mating trial system vs. Bailie's continuous mating trials system

2

3

4

1

Table 3. Break-even values of female calf price when the annualized net revenue in ONE* exceeds the revenue in CON* with culling in 6th parity

Conception rate	Annual discount rate (%)	BMS [†] number of culled cows			
		1	2	3	4
0.20	0	838	813	796	778
	5	878	859	846	833
	10	916	902	893	884
	Mean	877	858	845	832
0.40	0	729	712	700	689
	5	757	747	739	733
	10	785	780	777	776
	Mean	757	746	739	733
0.60	0	697	687	680	676
	5	729	728	726	728
	10	761	768	771	779
	Mean	729	728	726	728
0.80	0	682	680	677	678
	5	721	727	730	739
	10	759	774	783	800
	Mean	721	727	730	739

* One mating trial system vs. Bailie's continuous mating trials system

[†] Japanese Beef Marbling Standard number

2

1

Table 4. *Break-even values of female calf price when the annualized net revenue at culling in the 9th parity exceeds the revenue at culling in the 3rd parity for the continuous mating trial system*

Annual discount rate (%)	BMS* number of culled cows			
	1	2	3	4
0	794	947	1035	1164
5	828	990	1085	1221
10	861	1032	1133	1276
Mean	828	990	1084	1220

* Japanese Beef Marbling Standard number

2

- 1
- 1 Fig. 1. Quadratic regressions to estimate carcass price of a culled cow by culling
- 2 parity and Japanese Beef Marbling Standard number of the carcass.
- 3
- 4 Fig. 2. Scheme of the herd composition for Japanese beef cow-calf production.
- 5
- 6 Fig. 3. Sensitivity of (a) biological efficiency, (b) economic efficiency and (c)
- 7 annualized net revenue to changes in conception rate, weaning weight, weaning age,
- 8 calf market age, female calf price and annual discount rate in the one mating (left) and
- 9 continuous mating (right) trial systems.
- 10
- 11 Fig. 4. Effect of change in culling parity of cows on the biological efficiency in the
- 12 two mating trial systems.
- 13
- 14 Fig. 5. Effect of changes in culling parity, Japanese Beef Marbling Standard
- 15 number and female calf price on the economic efficiency in the two mating trial
- 16 systems.
- 17
- 18 Fig. 6. Effect of changes in culling parity, Japanese Beef Marbling Standard
- 19 number and female calf price on the annualized net revenue in the two mating trial
- 20 systems.
- 21
- 22 Fig. 7. Differences in optimal culling parity between economic efficiency and
- 23 annualized net revenue with changes in Japanese Beef Marbling Standard number,
- 24 female calf price, annual discount rate and the mating trial system.




Figure 1 Quadratic regressions to estimate carcass price of a culled cow by culling parity and Japanese Beef Marbling Standard number of the carcass.






Figure 2 Scheme of the herd composition for Japanese beef cow-calf production.


a) Change in Biological efficiency (Left: One mating, Right: Continuous mating)



b) Change in Economic efficiency (Left: One mating, Right: Continuous mating)




c) Change in Annualised net revenue (Left: One mating, Right: Continuous mating)



1
2 **Figure 3** Sensitivity of (a) biological efficiency, (b) economic efficiency and (c)
3 annualized net revenue to changes in conception rate, weaning weight, weaning
4 age, calf market age, female calf price and annual discount rate in the one mating
5 (left) and continuous mating (right) trial systems.


1



2 **Figure 4** Effect of change in culling parity of cows on the biological efficiency in
3 the two mating trial systems.


4

5



1
2 **Figure 5** Effect of changes in culling parity, Japanese Beef Marbling Standard
3 number and female calf price on the economic efficiency in the two mating trial
4 systems.


1



2

3 **Figure 6** Effect of changes in culling parity, Japanese Beef Marbling Standard
4 number and female calf price on the annualized net revenue in the two mating trial
5 systems.

1



2

3 **Figure 7** Differences in optimal culling parity between economic efficiency and
 4 annualized net revenue with changes in Japanese Beef Marbling Standard number,
 5 female calf price, annual discount rate and the mating trial system.

1
2
3

APPENDIX 1

Explanation of symbols and several fixed parameters in the model

Symbols	Units	Explanations	Fixed values
x (<i>subscript</i>)		m : males, f : females	
- Fixed parameters -			
t_{op}	d	Mean length of the oestrous cycle	21
t_{mfst}	d	Age at first mating	420
t_{preg}	d	Gestation length	285
$cmort$		Mortality rate before weaning	0.02
$mort$		Yearly mortality rate after weaning	0.02
t_{wean}	d	Weaning age	150
BW_x	kg	Birth weight (x 1.2 for males)	30 (females)
MW_x	kg	Mature weight (x 1.2 for males)	515 (females)
b		Wood's lactation curve parameter	0.073
c		Wood's lactation curve parameter	0.0056
TM	kg/year	Total annual milk yield	970
$kfmilk$		Efficiency of ME utilization of milk for growth	0.700
$kfsolid$		Efficiency of ME utilization of feed for growth	0.423
$milkME$	ME/kg	ME per kg DM of milk	5.12
$solidME$	ME/kg	ME per kg DM of dietary feed for calves	2.35
q		Overall metabolizability	0.60
q_c		Metabolizability of concentrates	0.70
q_r		Metabolizability of roughage	0.45
- Variables and functions -			
$WW_x(n)$	kg	Weaning weight (x 1.08 for males)	
$Cr(n)$		Conception rate of cows at parity n	
$T_{do}(n)$	d	Days open of cows at parity n	
$Mt(n)$	n	Average mating trial times of cows at parity n	
$T_{cl}(n)$	d	Period between parturition to next parturition	
$CP_x(pa)$	yen/kg	Carcass price of cows after parity at culling pa	
$NPV(pa)$	yen	Net present value	
$Day(pa)$	d	Days of age of cows with planned parity at culling	
$CF(i)$	yen	Daily cash flow	
$EDC(pa)$	yen	Equivalent dairy cash flow	
$AN(pa)$	yen	Annualized net revenue	
ydr		Annual discount rate	

(continued)		
t	d	Days of age
$dcmort$		Daily mortality of calves
$dmort$		Daily mortality of cows
A_x, B_x, K_x		Brody's growth curve parameters
$W_x(t)$	kg	Body weight
$DG_x(t)$	kg/d	Daily gain
t_c	d	Days from conception
$W_c(t_c)$	kg	Additional body weight for pregnant cow
a		Wood's lactation curve parameter
t_m	d	Days after calving
$Y(t_m)$	kg	Daily milk yield
$msrate(t)$		Ratio of dietary feed to total feed on a DM basis
$ME_{m_x}(t)$	Mcal/d	ME for maintenance
$NE_{g_x}(t)$	Mcal/d	NE for growth
$ME_{g_x}(t)$	Mcal/d	ME for growth
$ME_x(t)$	Mcal/d	Metabolizable energy intake
$memix(t)$	Mcal/kg	Integrated ME per kgDM for milk and dietary feed
$DML_x(t)$	kg/d	DM intake
k_f		Efficiency of ME utilization for growth
$E_{preg}(t_c)$	Mcal/d	Additional energy requirement for pregnancy
ME_{preg}	Mcal/d	Additional ME for pregnancy
kl		Efficiency of ME utilization for lactation
ME_{lac}	Mcal/d	Additional ME for lactation
$Cratcalf(t)$		Ratio of concentrates to dietary feed for calf
$CRate$		Ratio of concentrates to total feed
$N(pa, n)$	n	Number of cows at parity n
$rep(pa)$		Replacement rate for steady-state herd
$Tnb(pa)$	n	Total number of newborns
$Ncalf_x(pa, n)$	n	Number of newborns born from cows at n -th parity

APPENDIX 2

Mortality

Mortality of calves ($dcmort$) and cows ($dmort$) per day are calculated as:

$$dcmort = 1 - (1 - cmort)^{(1/t_{wean})}$$

$$dmort = 1 - (1 - mort)^{(1/365)}$$

where $cmort$ is proportional mortality before weaning, $mort$ is annual proportional mortality of cows and t_{wean} is days from birth to weaning.

Daily body weight change and daily milk yield

The form of the growth curve is expressed using birth weight (BW_x) and weaning weight (WW_x) as:

$$W_x(t) = (WW_x - BW_x)/t_{wean} \times t + BW_x \quad (t \leq t_{wean})$$

$$W_x(t) = A_x(1 - B_x e^{-K_x t}) \quad (t > t_{wean}) \text{ (kg)}$$

where A_x , B_x and K_x are Brody's growth curve parameters. From the functions, daily gain is expressed as:

$$DG_x(t) = (WW_x - BW_x)/t_{wean} \quad (t \leq t_{wean})$$

$$DG_x(t) = K_x(A_x - W_x(t)) \quad (t > t_{wean}) \text{ (kg/day)}.$$

Here, Brody's parameter A_x is assumed to be mature weight (MW_x). Since both functions of $DG_x(t)$ should be equal at weaning, parameter B_x and K_x can be calculated as:

$$B_x = (1 - WW_x/MW_x)e^{K_x t_{wean}}$$

$$K_x = ((WW_x - BW_x)/t_{wean})/(MW_x - WW_x).$$

These growth curve functions are modified with the change in weaning weight derived from the weaning weight change function (Eqn (1)) described in the text.

The total weight of the conceptus added to maternal weight for the last 2 months of pregnancy is estimated as (AFRC 1993):

$$W_c(t_c) = (BW/40) \times 10^{(2.932 - 3.347 \exp(-0.00406t_c))} \text{ (kg)}$$

where t_c is days from conception ($222 \leq t_c \leq t_{preg}$).

1 Daily milk yields of cows were estimated using Wood's lactation curve (Wood 1967) as:

$$2 \quad Y(t_m) = at_m^b e^{-ct_m} \quad (\text{kg/day})$$

3 where t_m is days after calving, a , b and c are Wood's parameters. In the model,
4 parameters b , c and total milk yield in lactation period (TM , kg) are given as animal traits
5 shown in Appendix 1. Using these parameters, the parameter a is calculated as:

$$6 \quad a = \left(\sum_{t_m}^{t_{\text{wean}}} t_m^b e^{-ct_m} \right) / TM.$$

7

8 *ME and DMI requirement*

9 The series of mathematical expressions about ME (Mcal/d) and DMI (kg/d) requirement of
10 Japanese Beef cow-calf production at an individual level was based on AFRC (1993) and
11 NARO (2009). Briefly, expressions to calculate ME and DMI are separated in two categories:
12 pre-weaning calves and post-weaning animals (steers, heifers and cows). ME calculations for
13 pre-weaning calves accounts for the energy from the cow's milk and dietary feed (roughage
14 and concentrates). The explanation of symbols is shown in Appendix 1.

15

16 [ME and DMI requirement for calves before weaning]

17

18 - Calculation of the ratio of dietary feed to total feed on a DM basis

$$19 \quad msrate(t) = 0 \quad (0 \leq t < 30)$$

$$20 \quad = 0.0004 \times 10^{-3} W_x(t)^3 - 0.0001 W_x(t)^2 + 0.0012 W_x(t) + 1.11 \quad (30 \leq t \leq 150)$$

21 - Calculation of ME requirement from milk and dietary feed

$$22 \quad MEm_x(t) = 0.1067 W_x(t)^{0.75}$$

$$23 \quad NEg_x(t) = (0.008 W_x(t) + 1.8) \times DG_x(t)$$

$$24 \quad MEg_x(t) = NEg_x(t) / (kfmilk \times msrate(t) + kfsolid \times (1 - msrate(t)))$$

$$25 \quad ME_x(t) = MEm_x(t) + MEg_x(t)$$

26 - Calculation of ME per kg DM of total feed using $msrate(t)$

$$1 \quad memix(t) = milkME \times msrate(t) + solidME \times (1 - msrate(t))$$

2 - Calculation of DMI from dietary feed

$$3 \quad DMI_x(t) = ME_x(t) / memix(t) \times ((solidME \times (1 - msrate(t))) / memix(t))$$

4

5 [ME and DMI for steers, heifers and cows after weaning]

6

7 - ME requirement for maintenance and growth

$$8 \quad MEm_x(t) = 0.1067W_x(t)^{0.75}$$

$$9 \quad NEg_x(t) = 0.0639W_x(t)^{0.75} \times DG_x(t)$$

$$10 \quad kf = 0.78q + 0.006$$

$$11 \quad MEg_x(t) = NEg_x(t) / kf$$

12 - For steers and non-pregnant heifers

$$13 \quad ME_x(t) = MEm_x(t) + MEg_x(t)$$

14 - For cows

$$15 \quad Epreg(t_c) = 1.542 \times t_c^{5.45601} \times 10^{-12}$$

$$16 \quad MEpreg = Epreg(t_c) / 0.15$$

$$17 \quad kl = 0.35q + 0.42$$

$$18 \quad MElac = 0.815 \times Y(t_m) / kl$$

$$19 \quad ME_x(t) = MEm_x(t) + MEg_x(t) + MEpreg + MElac$$

20 - Calculation of DMI

$$21 \quad DMI_x(t) = ME_x(t) / (4.4q)$$

22

23 *Feeding amounts of roughage and concentrates*

24

25 Quantity of roughage and concentrates for pre-weaning calves and post-weaning animals
26 were calculated as the product of DMI and the proportion of roughage or concentrates in the
27 total feed.

1 - For pre-weaning calves

2 The ratio of concentrates to dietary feed was estimated from the equation derived from JLTA
3 (2007) as:

$$4 \quad CRate_{calf}(t) = -0.003t + 0.9735 \quad (t \geq 30)$$

5 where $t \leq t_{wean}$. This ratio is variable with the age of calf (t).

6 - For post-weaning animals

7 The ratio of concentrates to total feed was estimated from the equation derived from Hirooka *et*
8 *al.* (1998a) as:

$$9 \quad CRate = ME_c / ME = (q - q_r) / (q_c - q_r)$$

10 where ME_c / ME is the proportion of concentrates, q_c and q_r are metabolizabilities of
11 concentrates and roughage, and q is the overall metabolizability of feeds. This ratio is not
12 variable when all parameters about metabolizability of feed are fixed.

13

APPENDIX 3

Herd composition dynamics

In the present study, the simulation was performed assuming that the number of replacement heifers at the start of simulation is 1.0. From this number, $N(pa, n)$ matrix was calculated and indicates the number of cows at reproduction time n when the cows are kept until culling parity pa ($n \leq pa$). The $N(pa, n)$ matrix is calculated with the mortality, and only in case of selecting one mating trial system described in the text, with the mortality and the conception rate as follows:

- In ONE

$$N(pa, 1) = (1.0 \times (1 - cmort) \times (1 - dmort)^{(t_{mfst} - t_{wean})}) \times Cr(1) \times (1 - dmort)^{t_{preg}}$$

$$N(pa, n) = (N(pa, n-1) \times (1 - dmort)^{(t_{cl} - t_{preg})}) \times Cr(n) \times (1 - dmort)^{t_{preg}}$$

where t_{cl} is a fixed calving interval (357 days).

- In CON

$$N(pa, 1) = 1.0 \times (1 - cmort) \times (1 - dmort)^{(T_{cl}(1) - t_{wean})}$$

$$N(pa, n) = N(pa, n-1) \times (1 - dmort)^{(T_{cl}(n))}$$

where $T_{cl}(n)$ is the calving interval function described in the text.

Using the matrix, total number of newborns ($Tnb(pa)$) and replacement rate of cows ($rep(pa)$) are expressed as:

$$Tnb(pa) = \sum_{i=1}^{pa} N(pa, i)$$

$$rep(pa) = 1.0 / (0.5 \times Tnb(pa)).$$

The denominator of the expression of $rep(pa)$ theoretically represents the sum of female calves when the sex ratio is 0.5. Using $N(pa, n)$ and $rep(pa)$, the numbers of newborn male calves ($Ncalf_m(pa, n)$) and non-replacement newborn female calves ($Ncalf_f(pa, n)$) born from cows at n -th parity are:

$$Ncalf_m(pa, n) = 0.5 \times N(pa, n)$$

$$Ncalf_f(pa, n) = (1 - rep(pa)) \times 0.5 \times N(pa, n)$$